arXiv:1407.5092v2 [math.CO] 8 Sep 2014

A Study on the Sparing Number of the Corona of Certain Graphs

K P Chithra [1,*], **K A Germina** [2] and **N K Sudev** [3]

[1]Naduvath Mana, Nandikkara P O
Thrissur - 680301, Kerala, India.
email: *chithrasudev@gmail.com*

[2]Department of Mathematics, School of Mathematical & Physical Sciences
Central University of Kerala, Kasaragod - 671316, Kerala, India.
email: *srgerminaka@gmail.com*

[3]Department of Mathematics, Vidya Academy of Science & Technology
Thalakkottukara, Thrissur - 680501, Kerala, India.
email: *sudevnk@gmail.com*

[*] Corresponding author.

Abstract

Let $\mathbb{N}_0$ be the set of all non-negative integers and $\mathcal{P}(\mathbb{N}_0)$ be its the power set. An integer additive set-indexer (IASI) is defined as an injective function $f : V(G) \to \mathcal{P}(\mathbb{N}_0)$ such that the induced function $f^+ : E(G) \to \mathcal{P}(\mathbb{N}_0)$ defined by $f^+(uv) = f(u)+f(v)$ is also injective, where $f(u)+f(v)$ is the sum set of $f(u)$ and $f(v)$. If $f^+(uv) = k \; \forall \; uv \in E(G)$, then f is said to be a k-uniform integer additive set-indexer. An integer additive set-indexer f is said to be a weak integer additive set-indexer if $|f^+(uv)| = \max(|f(u)|, |f(v)|) \; \forall \; uv \in E(G)$. We have some characteristics of the graphs which admit weak integer additive set-indexers. In this paper, we study about the sparing number of the corona of two graphs.

Key Words: Integer additive set-indexers, mono-indexed elements of a graphs, weak integer additive set-indexers, sparing number of a graph.
AMS Subject Classification: 05C78

1 Introduction

For all terms and definitions, not defined specifically in this paper, we refer to [6], [2] and [5]. Unless mentioned otherwise, all graphs considered here are simple, finite and have no isolated vertices.

Let $\mathbb{N}_0$ denote the set of all non-negative integers. For all $A, B \subseteq \mathbb{N}_0$, the sum of these sets is denoted by $A+B$ and is defined by $A+B = \{a+b : a \in A, b \in B\}$. The set $A + B$ is called the *sum set* of the sets A and B. If either A or B is countably infinite, then their sum set is also countably infinite. Hence, the sets we consider here are all finite sets of non-negative integers. The cardinality of a set A is denoted by $|A|$. The power set of a set A is denoted by $\mathcal{P}(A)$.

We define an integral multiple of a set A, as the set, denoted by $n.A$, every element of whose is an integral multiple of the corresponding element of A. Therefore, $|n.A| = |A|$.

Definition 1.1. [3] An *integer additive set-indexer* (IASI, in short) is defined as an injective function $f : V(G) \to \mathcal{P}(\mathbb{N}_0)$ such that the induced function $f^+ : E(G) \to \mathcal{P}(\mathbb{N}_0)$ defined by $f^+(uv) = f(u) + f(v)$ is also injective.

Lemma 1.2. *[4] If f is an integer additive set-indexer defined on a given graph G, then* $\max(|f(u)|, |f(v)|) \leq |f^+(uv)| \leq |f(u)|\,|f(v)|, \forall\ u, v \in V(G)$.

Definition 1.3. [4] An IASI f is said to be a *weak IASI* if $|f^+(uv)| = \max(|f(u)|, |f(v)|)$ for all $u, v \in V(G)$. A weak IASI f is said to be *weakly uniform IASI* if $|f^+(uv)| = k$, for all $u, v \in V(G)$ and for some positive integer k. A graph which admits a weak IASI may be called a *weak IASI graph*.

The following is the necessary and sufficient condition for a given graph to admit a weak IASI.

Lemma 1.4. *[4] A graph G admits a weak integer additive set-indexer if and only if every edge of G has at least one mono-indexed end vertex.*

Definition 1.5. [7] The cardinality of the labeling set of an element (vertex or edge) of a graph G is called the *set-indexing number* of that element. An element (a vertex or an edge) of graph which has the set-indexing number 1 is called a *mono-indexed element* of that graph.

Definition 1.6. [7] The *sparing number* of a graph G is defined to be the minimum number of mono-indexed edges required for G to admit a weak IASI and is denoted by $\varphi(G)$.

Theorem 1.7. *[7] A subgraph of weak IASI graph is also a weak IASI graph.*

Theorem 1.8. *[7] A graph G admits a weak IASI if and only if G is bipartite or it has at least one mono-indexed edge.*

Theorem 1.9. *[7] The sparing number of a bipartite graph G is* 0.

Theorem 1.10. *[7] An odd cycle C_n has a weak IASI if and only if it has at least one mono-indexed edge.*

Theorem 1.11. *[7] Let C_n be a cycle of length n which admits a weak IASI, for a positive integer n. Then, C_n has an odd number of mono-indexed edges when it is an odd cycle and has even number of mono-indexed edges, when it is an even cycle.*

Theorem 1.12. *[8] The graph $G_1 \cup G_2$ admits a weak IASI if and only if both G_1 and G_2 are weak IASI graphs.*

Theorem 1.13. *[7] The sparing number of a complete graph K_n is $\frac{1}{2}(n-1)(n-2)$.*

The admissibility of weak IASI by certain graph products is established in [9]. In this paper, our discussion is about the sparing number of a particular product, called corona, of two weak IASI graphs.

2 Corona of Weak IASI Graphs

Definition 2.1. [1] The *corona* of two graphs G_1 and G_2, denoted by $G_1 \odot G_2$, is the graph obtained by taking one copy of G_1 (which has n_1 vertices) and n_1 copies of G_2 and then joining the i-th point of G_1 to every point in the i-th copy of G_2. The number of vertices and edges in $G_1 \odot G_2$ are $n_1(1+n_2)$ and $m_1 + n_1m_2 + n_1n_2$ respectively, where n_i and m_i are respectively the order and size of the graph $G_i, i = 1, 2$.

In the corona $G_1 \odot G_2$ of given graphs G_1 and G_2, we take $|V(G_2)|$ layers or copies of G_2 and to establish the adjacency between all vertices of each copy to the corresponding vertex of G_1. The weak IASIs of G_1 and G_2 may not induce a weak IASI for $G_1 \odot G_2$. Hence, we have to define an IASI independently for a graph product.

The following theorem establishes a necessary and sufficient condition for the corona of two weak IASI graphs to admit a weak IASI.

Theorem 2.2. *[9] Let G_1 and G_2 be two weak IASI graphs on m and n vertices respectively. Then, $G_1 \odot G_2$ admits a weak IASI if and only if either G_1 is 1-uniform or it has r copies of G_2 that are 1-uniform, where r is the number of vertices in G_1 that are not mono-indexed.*

In view of Theorem 2.2, we examine the sparing number of the corona of certain graph structures. First, we consider the corona of two paths. The following result determines the sparing number $P_m \odot P_n$.

Theorem 2.3. *The sparing number of the corona $P_m \odot P_n$ is*

$$\varphi(P_m \odot P_n) = \begin{cases} \frac{1}{4}(m+1)(3n+1) & \text{if both } m \text{ and } n \text{ are odd} \\ \frac{3}{4}(m+1)n & \text{if both } m \text{ is odd and } n \text{ is even} \\ \frac{1}{4}(3mn+2n+m) & \text{if both } m \text{ is even and } n \text{ is odd} \\ \frac{1}{4}n(3m+4) & \text{if both } m \text{ and } n \text{ are even} \end{cases}$$

Proof. Let P_m and P_n be two paths of lengths m and n respectively. Note that P_m has $m+1$ vertices and P_n has $n+1$ vertices. Label the vertices of P_m alternately by distinct singleton sets and distinct non-singleton sets. Then, the number of mono-indexed edges in P_m is 0. Now, the number of mono-indexed vertices in P_m is $\frac{m+1}{2}$ if m is odd and $\frac{m}{2}$ if m is even. Hence, there are the following cases.

Case 1: Let m be odd. Then, exactly $\frac{m+1}{2}$ copies of P_n must be 1-uniform and the vertices of the remaining $\frac{m+1}{2}$ copies of P_n can be labeled alternately by distinct singleton and distinct non-singleton set-labels. In this context, we have the following subcases.

Subcase 1.1: Let n be an odd integer. Here, $\frac{m+1}{2}$ copies of P_n are 1-uniform. In the remaining $\frac{m+1}{2}$ copies, exactly $\frac{n+1}{2}$ vertices are mono-indexed. Then, $\frac{n+1}{2}$ edges between P_m and such a copy of P_n are mono-indexed. Therefore, the number of mono-indexed edges is $\frac{m+1}{2}n + \frac{(m+1)}{2}.\frac{(n+1)}{2} = \frac{1}{4}(m+1)(3n+1)$.

Subcase 1.2: Let n be an even integer. Here, $\frac{m+1}{2}$ copies of P_n are 1-uniform and in the remaining $\frac{m+1}{2}$ copies, $\frac{n}{2}$ vertices are mono-indexed. Therefore, $\frac{n}{2}$ edges between P_m and such a copy of P_n are mono-indexed. Hence, the number of mono-indexed edges is $\frac{m+1}{2}n + \frac{m+1}{2}.\frac{n}{2} = \frac{3}{4}(m+1)n$.

Case 2: Let m be even. Then, exactly $\frac{m}{2}$ copies of P_n must be 1-uniform and the vertices of the remaining $\frac{m+2}{2}$ copies of P_n can be labeled alternately by distinct singleton and distinct non-singleton set-labels. Here, we have the following subcases.

Subcase 2.1: Let n be an odd integer. Here, $\frac{m}{2}$ copies of P_n are 1-uniform. In the remaining $\frac{m+2}{2}$ copies, exactly $\frac{n+1}{2}$ vertices are mono-indexed. Then, $\frac{n+1}{2}$ edges between P_m and such a copy of P_n are mono-indexed. Therefore, the number of mono-indexed edges is $\frac{m}{2}n + \frac{(m+2)}{2}.\frac{(n+1)}{2} = \frac{1}{4}(3mn+2n+m)$.

Subcase 2.2: Let n be an even integer. Here, $\frac{m+2}{2}$ copies of P_n are 1-uniform and in the remaining $\frac{m}{2}$ copies, $\frac{n}{2}$ vertices are mono-indexed. Therefore, $\frac{n}{2}$ edges between P_m and such a copy of P_n are mono-indexed. Therefore, the number of mono-indexed edges is $\frac{m+2}{2}n + \frac{m}{2}.\frac{n}{2} = \frac{1}{4}n(3m+4)$. □

Next, we discuss the corona of two graphs in which one is a cycle. First, recall the following remark.

Remark 2.4. [7] Due to Theorem 1.10, we observe that the number of mono-indexed vertices in the cycle C_n of length n is at least $\frac{(n+1)}{2}$ if n is odd and is at least $\frac{n}{2}$, if n is even.

By Theorem 2.2, The corona of a cycle C_m and a path P_n, denoted by $C_m \odot P_n$, admits a weak IASI if and only if either C_m is 1-uniform or r copies of P_n that are 1-uniform, where r is the number of vertices of C_m that are not mono-indexed and $P_n \odot C_m$ admits a weak IASI if and only if either P_n is 1-uniform or r copies of C_m are 1-uniform, where r is the number of vertices of P_n that are not mono-indexed. Hence, we have

Theorem 2.5. *The sparing number of $C_m \odot P_n$ is*

$$\varphi(C_m \odot P_n) = \begin{cases} \frac{3}{4}mn & \text{if both } m \text{ and } n \text{ are odd} \\ \frac{1}{4}m(3n+1) & \text{if both } m \text{ is odd and } n \text{ is even} \\ 1+\frac{1}{4}m(3m-1) & \text{if both } m \text{ is even and } n \text{ is odd} \\ \frac{1}{4}[n(3m+1)+(m+5)] & \text{if both } m \text{ and } n \text{ are even} \end{cases}$$

Proof. Consider a cycle C_m and a path P_n. Note that the path P_n has $m+n$ vertices, while C_m has m vertices. Here, we consider the following cases.

Case-1: Let m is even. Then, exactly $\frac{m}{2}$ vertices of C_m have non-singleton set-label. Therefore, by Theorem 2.2, $C_m \odot P_n$ contains $\frac{m}{2}$ copies of P_n that are 1-uniform. The remaining $\frac{m}{2}$ copies of P_n can be labeled alternately by distinct singleton and non-singleton sets. For these $\frac{m}{2}$ copies, we have the following subcases.

Subcase-1.1: Let n be an even integer. Then, $\frac{n+2}{2}$ vertices can be labeled by non-singleton set-labels and the remaining $\frac{n}{2}$ vertices have singleton set-labels. Therefore, there are $\frac{n}{2}$ mono-indexed edges between C_m and these copies of P_n. Hence, the total number of mono-indexed edges in $C_m \odot P_n$ is $\frac{m}{2}n + \frac{m}{2}\frac{n}{2} = \frac{3}{4}mn$.

Subcase-1.2: Let n be an odd integer. Then, for each copy of P_n corresponding to a mono-indexed vertex of C_m, there are $\frac{n+1}{2}$ vertices that can be labeled by non-singleton sets and $\frac{n+1}{2}$ vertices can be labeled by singleton sets. Therefore, there are $\frac{n+1}{2}$ mono-indexed edges between C_m and these copies of P_n. Hence, the total number of mono-indexed edges in $C_m \odot P_n$ is $\frac{m}{2}n + \frac{m}{2}.\frac{n+1}{2} = \frac{1}{4}m(3n+1)$.

Case-2: Let m is odd. Then, by Theorem 1.10, the cycle C_m has at least one mono-indexed edge. That is, C_m has $\frac{m-1}{2}$ vertices that are not mono-indexed and $\frac{m+1}{2}$ mono-indexed vertices. Then, $\frac{m-1}{2}$ copies of P_n are 1-uniform and the vertices of the corresponding copies of remaining $\frac{m+1}{2}$ vertices of C_m, can be labeled alternately by singleton sets and non-singleton sets. In this case also, we have two options.

Subcase-2.1: Let n be an even integer. Then, as explained in the above case, $\frac{n}{2}$ vertices have singleton set-labels and the remaining $\frac{n+2}{2}$ vertices can be labeled by non-singleton set-labels. Therefore, there are $\frac{n}{2}$ mono-indexed edges between C_m and these copies of P_n. Hence, the total number of mono-indexed edges in $C_m \odot P_n$ is $1 + \frac{m-1}{2}n + \frac{m+1}{2}.\frac{n}{2} = 1 + \frac{1}{4}n(3m-1)$.

Subcase-2.2: Let n be an odd integer. Then, for each copy of P_n corresponding to a mono-indexed vertex of C_m, there are $\frac{n+1}{2}$ vertices that can be mono-indexed and the remaining $\frac{n+1}{2}$ vertices can be labeled by non-singleton sets. Therefore, there are $\frac{n+1}{2}$ mono-indexed edges between C_m and these copies of P_n. Hence, the total number of mono-indexed edges in $C_m \odot P_n$ is $1 + \frac{m-1}{2}n + \frac{m+1}{2}.\frac{n+1}{2} = \frac{1}{4}[n(3m+1)+(m+5)]$ □

Since $C_m \odot P_n$ and $P_n \odot C_m$ are non-isomorphic graphs, the sparing numbers of these graphs are also different. Therefore, we prove the following theorem for the corona $P_n \odot C_m$.

Theorem 2.6. *The sparing number of $P_n \odot C_m$ is*

$$\varphi(P_n \odot C_m) = \begin{cases} \frac{3}{4}(m+1)(n+1) & \text{if both } m \text{ and } n \text{ are odd} \\ \frac{3}{4}m(n+1) & \text{if both } m \text{ is even and } n \text{ is odd} \\ \frac{1}{4}[3n(m+1)+2(m+3)] & \text{if both } m \text{ is odd and } n \text{ is even} \\ \frac{1}{4}m(3n+2) & \text{if both } m \text{ and } n \text{ are even.} \end{cases}$$

Proof. We consider the following cases.

Case-1: Let n is odd. then, $\frac{n+1}{2}$ vertices can be labeled by non-singleton sets and $\frac{n+1}{2}$ vertices can be labeled by singleton sets. Therefore, by Theorem 2.2, $\frac{n+1}{2}$ copies of C_m must be 1-uniform. For the remaining $\frac{n+1}{2}$ copies of C_m, the vertices can be labeled alternately by singleton sets and non-singleton sets respectively. In this context, we have the following subcases.

Subcase-1.1: Let m be an odd integer. Then, each copy C_m, corresponding to the mono-indexed vertices of P_n, can have at least one mono-indexed edge. In this case, it has $\frac{m+1}{2}$ mono-indexed vertices and $\frac{m-1}{2}$ vertices that are not mono-indexed. That is, there exist $\frac{m+1}{2}$ edges between P_n and each of the relevant copies of C_m. Hence, the total number of mono-indexed edges in $P_n \odot C_m$ is $\frac{n+1}{2}m + \frac{n+1}{2}.\frac{m+1}{2} + \frac{n+1}{2} = \frac{3}{4}(m+1)(n+1)$.

Subcase-1.2: Let m be an even integer. Then, $\frac{m}{2}$ vertices can be labeled by singleton set-labels and the remaining $\frac{m}{2}$ vertices have non-singleton set-labels. Then, there exist $\frac{m}{2}$ edges between P_n and each of the relevant copies of C_m. Hence, the total number of mono-indexed edges in $P_n \odot C_m$ is $\frac{n+1}{2}m + \frac{n+1}{2}\frac{m}{2} = \frac{3}{4}m(+1)n$.

Case-2: Let n is even. Then, $\frac{n+2}{2}$ vertices of P_n have singleton set-label and $\frac{n}{2}$ vertices of P_n have non-singleton set-label. Therefore, by Theorem 2.2, $\frac{n+2}{2}$ copies of C_m are 1-uniform in $C_m \odot P_n$ and the remaining $\frac{n}{2}$ copies of C_m can be labeled alternately by distinct singleton and non-singleton sets. For these $\frac{m}{2}$ copies, we have the following subcases.

Subcase-2.1: Let m be an odd integer. Then, for each copy of C_m corresponding to a mono-indexed vertex of P_n, there are $\frac{m+1}{2}$ vertices that can be mono-indexed and the remaining $\frac{m-1}{2}$ vertices can be labeled by non-singleton sets. Then, there exist $\frac{m+1}{2}$ edges between P_n and each of the relevant copies of C_m. Hence, the total number of mono-indexed edges in $P_n \odot C_m$ is $\frac{n}{2}m + \frac{n+2}{2}.\frac{m+1}{2} + \frac{n+2}{2} = \frac{1}{4}[3n(m+1) + 2(m+3)]$.

Subcase-2.2: Let m be an even integer. Then, $\frac{m}{2}$ vertices have singleton set-labels and the remaining $\frac{m}{2}$ vertices can be labeled by non-singleton set-labels. Then, there exist $\frac{m}{2}$ edges between P_n and each of the relevant copies of C_m. Therefore, the total number of mono-indexed edges in $P_n \odot C_m$ is $\frac{n}{2}m + \frac{n+2}{2}.\frac{m}{2} = \frac{1}{4}m(3n+2)$. □

We now discuss on the the sparing number of the corona of two cycles. Let C_m and C_n be two cycles that admit weak IASI. Then, by Theorem 2.2, the corona $C_m \odot C_n$ admits a weak IASI if and only if C_m is 1-uniform or r copies of C_m are 1-uniform, where r is the number of vertices in C_m that are not mono-indexed. Hence, we have

Theorem 2.7. *The sparing number of $C_m \odot C_n$ is*

$$\varphi(C_m \odot C_n) = \begin{cases} \frac{3}{4}mn & \text{if } m \text{ and } n \text{ are even} \\ \frac{3}{4}m(n+1) & \text{if } m \text{ is even and } n \text{ is odd} \\ 1 + \frac{1}{4}(3m-1)n & \text{if } m \text{ is odd and } n \text{ is even} \\ 2 + \frac{1}{4}(3m-1)(n+1) & \text{if } m \text{ and } n \text{ are odd.} \end{cases}$$

Proof. Let C_m and C_n be two cycles which admit weak IASIs. Let $G = C_m \odot C_n$. We consider the problem in two cases.

Case-1: Let m be even. Then, C_m has $\frac{m}{2}$ mono-indexed vertices and has $\frac{m}{2}$ vertices that are not mono-indexed. Then, by Theorem 2.2, $\frac{m}{2}$ copies of C_n are 1-uniform. Let C_{n_i} be the copy of C_n corresponding to the i-th vertex of C_m. Let $\mathbb{C}$ be the set of all copies C_{n_i} in G that are 1-uniform and $\mathbb{C}'$ be the set of all copies C_{n_i} in G that are not 1-uniform. The number of mono-indexed edges in in $\mathbb{C}$ is $\frac{mn}{2}$. To find the number of other mono-indexed edges in G, we consider the following subcases.

Subcase-1.1: Let n is even. Then, each C_{n_i} in $\mathbb{C}'$ has (at least) $\frac{n}{2}$ mono-indexed vertices and need not have any mono-indexed edge. Since the corresponding vertex v_i in C_1 is mono-indexed, the number of mono-indexed edges in the component $C_{n_i} + \{v_i\}$ is $\frac{n}{2}$. Therefore, the number of mono-indexed edges connecting the cycle C_m and the elements of $\mathbb{C}'$ is $\frac{m}{2}.\frac{n}{2} = \frac{mn}{4}$. Hence, the total number of mono-indexed edges in G is $\frac{mn}{2} + \frac{mn}{4} = \frac{3}{4}mn$.

Subcase-1.2: Let n is odd. Then, each C_{n_i} in $\mathbb{C}'$ has (at least) $\frac{n+1}{2}$ mono-indexed vertices and has (at least) one mono-indexed edge. Therefore, the total number of mono-indexed edges in $\mathbb{C}'$ is $\frac{m}{2}$. Now, the number of mono-indexed edges in $C_{n_i} + \{v_i\}$ is at least $\frac{(n+1)}{2}$. Therefore, the number of mono-indexed edge connecting the cycle C_m and the elements of $\mathbb{C}'$ is $\frac{m}{2}.\frac{(n+1)}{2}$. Hence, the total number of mono-indexed edges in G is $\frac{mn}{2} + \frac{m}{2}.\frac{(n+1)}{2} + \frac{m}{2} = \frac{3}{4}m(n+1)$.

Case-2: Let m be odd. Then, C_m has $\frac{(m+1)}{2}$ mono-indexed vertices and has $\frac{(m-1)}{2}$ vertices that are not mono-indexed. Also, C_m must have (at least) one mono-indexed edge. Then, by Theorem 2.2, $\frac{(m-1)}{2}$ copies of C_n are 1-uniform. The number of mono-indexed edges in $\mathbb{C}$ is (at least) $\frac{(m-1)n}{2}$. To find the number of other mono-indexed edges in G, we consider the following subcases.

Subcase-1.1: Let n is even. Then, each C_{n_i} in $\mathbb{C}'$ has (at least) $\frac{n}{2}$ mono-indexed vertices and need not have any mono-indexed edge. Since the corresponding vertex v_i in C_m is mono-indexed, the number of mono-indexed edges in the component $C_{n_i} + \{v_i\}$ is at least $\frac{n}{2}$. Therefore, the number of mono-indexed edge connecting the cycle C_1 and the elements of $\mathbb{C}'$ is at least $\frac{(m+1)}{2}.\frac{n}{2} = \frac{(m+1)n}{4}$. Hence, the total number of mono-indexed edges in G is $1 + \frac{(m-1)n}{2} + \frac{(m+1)n}{4} = 1 + \frac{1}{4}(3m-1)n$.

Subcase-1.2: Let n is odd. Then, each C_{n_i} in $\mathbb{C}'$ has (at least) $\frac{n+1}{2}$ mono-indexed vertices and has (at least) one mono-indexed edge. Therefore, the total number of mono-indexed edges in $\mathbb{C}'$ is at least $\frac{(m+1)}{2}$. Now, the number of mono-indexed edges in $C_{n_i} + \{v_i\}$ is at least $\frac{(n+1)}{2}$. Therefore, the number of mono-indexed edge connecting the cycle C_m and the elements of at least $\mathbb{C}'$ is $\frac{(m+1)}{2}.\frac{(n+1)}{2}$. Hence, the total number of mono-indexed edges in G is at least $1 + \frac{(m-1)n}{2} + \frac{(m+1)(n+1)}{4} + \frac{(m+1)}{2} = 2 + \frac{1}{4}(3m-1)(n+1)$. □

We now proceed to discuss about the sparing numbers of the corona of two graphs, at least one of them being a complete graph. In the following theorem, we estimate the sparing number of the corona of two complete graphs.

Theorem 2.8. *The sparing number of the corona $K_m \odot K_n$ of two complete graphs K_m and K_n is $\frac{1}{2}[(m-1)(m-2) + mn(n-1)]$.*

Proof. The number of edges in $K_m \odot K_n$ is $|E| = \frac{1}{2}m(m-1) + mn + \frac{1}{2}mn(n-1) = \frac{1}{2}m[n^2 + n - 1 + m]$.

By Theorem 1.13, the graph K_m has at most one vertex that is not mono-indexed. Hence, in $K_m \odot K_n$, only one copy of K_n, say K_{n1}, needs to be 1-uniform. Therefore, all the n edges connecting K_m and the copy K_{n1} are not mono-indexed. All other $(m-1)$ copies of K_n can have one vertex having non-singleton set-label.

That is, all copies of K_n, except K_{n1}, have $(n-1)$ edges that are not mono-indexed. Therefore, the total number of edges in $K_m \odot K_n$ that are not mono-indexed , is $(m-1)+n+(m-1)n = mn+m-1$.

Hence, the number of mono-indexed edges in $K_m \odot K_n$ is $\varphi(K_m \odot K_n) = \frac{1}{2}m[n^2+n-1+m] - (mn+m-1) = \frac{1}{2}[(m-1)(m-2)+mn(n-1)]$. □

Next, we consider the corona of a path and a complete graph. We determine the sparing number of $P_m \odot K_n$ in the following theorem.

Theorem 2.9. *The sparing number of $P_m \odot K_n$ is $\varphi(P_m \odot K_n) = \frac{1}{2}n(n-1)(m+1)$.*

Proof. Consider the following cases.

Case-1: If m is even, Then P_m has $\frac{m}{2}$ vertices that are not mono-indexed and $\frac{m+2}{2}$ mono-indexed vertices. Therefore, $\frac{m}{2}$ copies of K_n must be 1-uniform. For the remaining copies of K_n, one vertex can have a mono-indexed set label and hence, there are $(n-1)$ mono-indexed edges between P_m and these copies of K_n. Therefore, the total number of mono-indexed edges in $P_m \odot K_n$ is $\frac{m}{2}\frac{n(n-1)}{2} + \frac{m+2}{2}\frac{(n-1)(n-2)}{2} + \frac{m+2}{2}(n-1) = \frac{1}{2}n(n-1)(m+1)$.

Case-2: If m is odd, Then, there are P_m $\frac{m+1}{2}$ mono-indexed vertices and P_m $\frac{m+1}{2}$ vertices that are not mono-indexed. Therefore, $\frac{m+1}{2}$ copies of K_n must be 1-uniform. For the remaining copies of K_n, there are $(n-1)$ mono-indexed edges between P_m and these copies of K_n. Therefore, the total number of mono-indexed edges in $P_m \odot K_n$ is $\frac{m+1}{2}\frac{n(n-1)}{2} + \frac{m+1}{2}\frac{(n-1)(n-2)}{2} + \frac{m+1}{2}(n-1) = \frac{1}{2}n(n-1)(m+1)$.

This completes the proof. □

The following Theorem estimates the sparing number of $K_n \odot P_m$.

Theorem 2.10. *The sparing number of $K_n \odot P_m$ is*

$$\varphi(K_n \odot P_m) = \begin{cases} \frac{1}{2}[(n-1)^2 + m(n+1)] & \text{if } m \text{ is odd} \\ \frac{1}{2}[(n-1)(n-2) + m(n+1) & \text{if } m \text{ is even.} \end{cases}$$

Proof. By Theorem 1.13, the complete graph K_n has $\frac{1}{2}(n-1)(n-2)$ mono-indexed edges. Since K_n has only one vertex that is not mono-indexed, only one copy of P_m needs to be 1-uniform and the remaining $(n-1)$ copies of P_m can be labeled alternately by singleton and non-singleton sets. For these copies of P_m we have the following cases.

Case-1: If m is odd, then each copy of P_m has $\frac{m+1}{2}$ mono-indexed vertices and hence the total number of mono-indexed edges is $\frac{(n-1)(n-2)}{2} + \frac{(n-1)(m+1)}{2} + m = \frac{1}{2}[(n-1)^2 + m(n+1)]$.

Case-2: If m is even, then each copy of P_m has $\frac{m}{2}$ mono-indexed vertices and $\frac{m+2}{2}$ vertices that are not mono-indexed. Therefore, the total number of mono-indexed edges $k_n \odot P_m$ is $\frac{(n-1)(n-2)}{2} + \frac{n-1}{2}m + m = \frac{1}{2}[(n-1)(n-2) + m(n+1)]$. □

The following theorems estimate the sparing number of the coronas of two graphs in which one is a cycle and the other is a complete graph.

Theorem 2.11. *The sparing number of* $C_m \odot K_n$ *is* $\varphi(P_m \odot K_n) = \frac{1}{2}mn(n-1)$.

Proof. Consider the following cases.

Case-1: If m is odd, Then C_m has $\frac{m-1}{2}$ vertices that are not mono-indexed and $\frac{m+1}{2}$ mono-indexed vertices. Therefore, $\frac{m-1}{2}$ copies of K_n must be 1-uniform. For the remaining copies of K_n, one vertex can have a mono-indexed set label and hence, there are $(n-1)$ mono-indexed edges between C_m and these copies of K_n. Therefore, the total number of mono-indexed edges in $C_m \odot K_n$ is $\frac{m-1}{2}\frac{n(n-1)}{2} + \frac{m+1}{2}\frac{(n-1)(n-2)}{2} + \frac{m+1}{2}(n-1) = \frac{1}{2}mn(n-1)$.

Case-2: If m is even, Then, there are $\frac{m}{2}$ mono-indexed vertices and $\frac{m}{2}$ vertices that are not mono-indexed. Therefore, $\frac{m}{2}$ copies of K_n must be 1-uniform. For the remaining copies of K_n, there are $(n-1)$ mono-indexed edges between C_m and these copies of K_n. Therefore, the total number of mono-indexed edges in $C_m \odot K_n$ is $\frac{m}{2}\frac{n(n-1)}{2} + \frac{m}{2}\frac{(n-1)(n-2)}{2} + \frac{m}{2}(n-1) = \frac{1}{2}mn(n-1)$. □

Theorem 2.12. *The sparing number of* $K_n \odot C_m$ *is*

$$\varphi(K_n \odot C_m) = \begin{cases} \frac{1}{2}[(n-1)^2 + m(n+1)] & \text{if } m \text{ is odd} \\ \frac{1}{2}[(n-1)(n-2) + m(n+1) & \text{if } m \text{ is even.} \end{cases}$$

Proof. By Theorem 1.13, the complete graph K_n has $\frac{1}{2}(n-1)(n-2)$ mono-indexed edges. Since K_n has only one vertex that is not mono-indexed, only one copy of C_m needs to be 1-uniform and the remaining $(n-1)$ copies of C_m can be labeled alternately by singleton and non-singleton sets. For these copies of C_m we have the following cases.

Case-1: If m is odd, then each copy of C_m has $\frac{m+1}{2}$ mono-indexed vertices and $\frac{m-1}{2}$ vertices that are not mono-indexed. Hence the total number of mono-indexed edges in $K_n \odot C_m$ is $\frac{(n-1)(n-2)}{2} + \frac{(n-1)(m+1)}{2} + m = \frac{1}{2}[(n-1)^2 + m(n+1)]$.

Case-2: If m is even, then each copy of P_m has $\frac{m}{2}$ mono-indexed vertices and $\frac{m}{2}$ vertices that are not mono-indexed. Therefore, the total number of mono-indexed edges in $K_n \odot C_m$ is $\frac{(n-1)(n-2)}{2} + \frac{n-1}{2}m + m = \frac{1}{2}[(n-1)(n-2) + m(n+1)]$. □

Next, we study about the sparing number of the corona of two graphs at least one of which is a bipartite graph. In the following theorem, we verify the corona of two complete bipartite graphs.

Theorem 2.13. *Let* $G_1 = K_{m_1,n_1}$ *and* $G_2 = K_{m_2,n_2}$ *be two complete bipartite graphs, where* $m_i \leq n_i$ *for* $i = 1, 2$. *Then, the sparing number of* $G_1 \odot G_2$ *is* $m_2(m_1n_2 + n_1)$.

Proof. For $i = 1, 2$, let $G_i(U_i, V_i, E_i)$ be a bipartite graph with $|U_i| = m_i$, $|V_i| = n_i$ and $|E_i = m_in_i$, where $m_i \leq n_i$. By Theorem 1.9, $\varphi(G_i) = 0$.

Label the vertices in U_1 of G_1 by distinct non-singleton sets and the vertices of V_1 of G_1 by distinct singleton sets. Label the vertices in U_2 of G_2 by distinct singleton sets and the vertices of V_2 of G_2 by distinct non-singleton sets, which are distinct from the set-labels of G_1.

Now, take $|V(G_1)|$ copies of G_2 and draw edges between the all vertices of each copy of G_2 and the corresponding vertex of G_1. Then, all the copies of G_2

corresponding to the vertices in U_1 of G_1 must be 1-uniform. That is, m_1 copies of G_2 are 1-uniform in $G_1 \odot G_2$. Therefore, the total number of mono-indexed edges among these copies of G_2 is $m_1 m_2 n_2$.

Label the remaining n_1 copies of G_2 in such a way that the set-labels of the vertices of a copy of G_2 are the (suitable) integral multiple of the set-labels of the corresponding set-labels of G_2 in such a way that no two vertices in G_1 and all copies of G_2 have a same set-label. then, the number edges between a vertex of G_1 (which are in V_1), and the vertices of the corresponding copy of G_2 is m_2. Therefore, the number of such mono-indexed edges among the vertices in V_1 of G_1 and the corresponding copies of G_2 is $n_1 m_2$.

Therefore, the total number of mono-indexed edges in $G_1 \odot G_2$ is $m_1 m_2 n_2 + n_1 m_2 = m_2(m_1 n_2 + n_1)$. □

Theorem can be generalised for all bipartite graphs as follows and the proof of the theorem is similar to the proof of Theorem 2.13.

Theorem 2.14. *Let $G_1(U_1, V_1, E_1)$ and $G_2(U_2, V_2, E_2)$ be two bipartite graphs, where $|U_i| \leq |V_i|$ for $i = 1, 2$. Then, the sparing number of $G_1 \odot G_2$ is $|U_1|\,|E_2| + |V_1||U_2|$.*

Now that we have estimated the sparing number of the corona of two bipartite graphs, we need to find out the sparing number of the corona of a bipartite graph and an odd cycle.

Theorem 2.15. *Let C_n be an odd cycle and $G(V_1, V_2, E)$ be a bipartite graph, where $|V_1| = r$, $|V_2| = s$ and $|E| = q$. Then,*

$$\begin{aligned} \varphi(C_n \odot G) &= \frac{1}{2}[(n-1)q + (n+1)r] \\ \text{and} \quad \varphi(G \odot C_n) &= \frac{1}{2}[2rn + s(n+1)]. \end{aligned}$$

Proof. Without loss of generality, let $r \leq s$. In G, label all the vertices in V_1 by distinct singleton sets and label all the vertices in V_1 by distinct non-singleton sets.
Case-1: Since C_n is an odd cycle, it has $\frac{n+1}{2}$ mono-indexed vertices and $\frac{n-1}{2}$ vertices that are not mono-indexed. Then, $\frac{n-1}{2}$ copies of G, must be 1-uniform in $C_n \odot G$. Therefore, the total number of mono-indexed edges in all these copies is $\frac{n-1}{2}q$. Now, label the vertices of the remaining copies of G, by suitable integral multiples of the set-labels of corresponding vertices of G in such a way that no two vertices in $C_n \odot G$ have the same set-label. Then, no two vertex in these copies are mono-indexed. But, there are r edges connecting a mono-indexed vertex of C_n and the corresponding copy of G. The total number of such mono-indexed edges is $\frac{n+1}{2}r$. Therefore, the sparing number of $C_n \odot G$ is $\frac{1}{2}[(n-1)q + (n+1)r]$.
Case-2: Since $r \leq S$, label all the vertices in V_1 of G by distinct non-singleton sets and label all the vertices in V_2 by distinct singleton sets. Then, r copies of C_n must be 1-uniform in $G \odot C_n$. The number mono-indexed copies all together is rn. Now, the remaining copies of C_n, corresponding to the vertices in V_2 of G, contain $\frac{n+1}{2}$ mono-indexed vertices each and hence there are $\frac{n+1}{2}$ mono-indexed edges between a vertex in V_2 and its corresponding copy of C_n is $\frac{n+1}{2}$. Therefore,

the number of such mono-indexed edges is $\frac{n+1}{2}s$. Hence, the sparing number of $G \odot C_n$ is $rn + \frac{n+1}{2}s = \frac{1}{2}[2rn + (n+1)s]$. □

It remains to find the sparing number of the corona of two graphs, one of which is a complete graph and the other is a bipartite graph. Then, we have

Theorem 2.16. *Let K_n be an odd cycle and $G(V_1, V_2, E)$ be a bipartite graph, where $|V_1| = r$, $|V_2| = s$ and $|E| = q$. Then,*

$$\begin{aligned} \varphi(K_n \odot G) &= \frac{1}{2}[(n-1)(n-2) + 2q + 2r(n-1)] \\ \textit{and} \quad \varphi(G \odot K_n) &= \frac{1}{2}(n-1)[rn + s(n-2)]. \end{aligned}$$

Proof. Let $r \leq s$. In G, label all the vertices in V_1 by distinct singleton sets and label all the vertices in V_1 by distinct non-singleton sets. In G, label all the vertices in V_1 by distinct singleton sets and label all the vertices in V_1 by distinct non-singleton sets.

Case-1: in K_n, every vertex is adjacent to all other vertices of it, only one vertex in K_n can have a non-singleton set-label. Therefore, one copy of G must be 1-uniform in $K_n \odot G$. The vertices of the remaining copies of G shall be labeled by the integral multiples of the set-labels of the corresponding vertices of G. Then, the total number of mono-indexed edges in $K_n \odot G$ is $q + (n-1)r + \frac{1}{2}(n-1)(n-2) = \frac{1}{2}[(n-1)(n-2) + 2q + 2r(n-1)]$.

Case-2: In $G \odot K_n$, r copies of K_n must be 1-uniform. The remaining copies of K_n can have $(n-1)$ vertices that are mono-indexed. Therefore, the total number of mono-indexed edges is $r\frac{1}{2}n(n-1) + s\frac{1}{2}(n-1)(n-2) = \frac{1}{2}(n-1)[rn + s(n-2)]$.

□

3 Conclusion

In this paper, we have discussed about the sparing number of the corona of weak IASI graphs. Some problems in this area are still open. Uncertainty in the adjacency pattern of different graphs makes this study complex. An investigation to verify the admissibility of weak IASIs by other graph products of two arbitrary graphs and to determine the corresponding sparing numbers seems to be fruitful.

References

[1] R Frucht and F Harary (1970). *On the Corona of Two Graphs*, Aequationes Math., **4**(3), 322-325.

[2] J A Gallian, (2011). *A Dynamic Survey of Graph Labelling*, The Electronic Journal of Combinatorics (DS 16).

[3] K A Germina and T M K Anandavally, (2012). *Integer Additive Set-Indexers of a Graph:Sum Square Graphs*, Journal of Combinatorics, Information and System Sciences, **37**(2-4), 345-358.

[4] K A Germina and N K Sudev, (2013). *On Weakly Uniform Integer Additive Set-Indexers of Graphs*, Int. Math. Forum, **8**(37), 1827-1834.

[5] R Hammack, W Imrich and S Klavzar (2011). **Handbook of Product graphs**, CRC Press.

[6] F Harary, (1994). **Graph Theory**, Addison-Wesley Publishing Company Inc.

[7] N K Sudev and K A Germina, (2014). *A Characterisation of Weak Integer Additive Set-Indexers of Graphs*, ISPACS J. Fuzzy Set Valued Analysis, **2014**, Article Id: jfsva-0189, 7 pages.

[8] N K Sudev and K A Germina, (2014). *Weak Integer Additive Set-Indexers of Graph Operations*, Global J. Math. Sciences: Theory and Practical, **6**(1),25-36.

[9] N K Sudev and K A Germina, *Weak integer Additive Set-Indexers of Certain Graph Products*, submitted.